3. Purification of community water supplies by flocculation, Filtration and adequate chlorination.

c) Control of susceptible population :

Vaccines: Several inactivated or live attenuated vaccines against Hepatitis A is available to protect against the infection.


Treatment

1. The patient has to be provided with adequate rest.
2. Bland diet should be provided
3. Avoid Alcohol

Acute Diarrheal Diseases

Definition

According to WHO Acute diarrhea is defined as an abnormally frequent discharge of semisolid or fluid faecal matter from the bowel, lasting less than the 14 days by WHO.



Causative Organism

Bacteria: Escherichia coli, Shigella, salmonella etc.,

Virus: Rota virus, adenovirus etc.

Parasites: Endamoeba hystelytica, Giardia lamblin etc.

Mode Of Transmission - Direct transmission - Faeco – oral route

Incubation Period - Few Hour to one day.

Clinical Manifestation

1. Sunken eyes
2. Tachycardia
3. Hypotension
4. Irritable and restlessness
5. Pallor
6. Rapid respiration
7. Sudden collapse if not treated properly
8. Stools loose and fluid in consistency, greenish or yellow green in colour, may contain mucus or blood.
9. Vomiting
10. Fever
11. Poor skin turgor, dry skin and dry mouth
12. Sunken fontanels in children

Diagnosis – Stool test

Complications

1. Persistent diarrhea
2. Malnutrition
3. Vitamins and mineral deficiencies
4. Hypoglycemia resulting in convulsions and brain damage
5. Electrolyte loss
6. Hypovolemic shock
7. Acute renal failure

Control Measures

1. Promote exclusive breastfeeding
2. Immunization
3. Using sanitary latrines
4. Keep food and water clean and closed.
5. Wash hands before eating and after defaecation.

Treatment

Oral Rehydration Therapy: Give some available liquids like rice water, oral rehydration solution (ORS) packet to be dissolved in one litre of drinking water and stir with clean spoon, till it dissolves. Give ¼ to ½ cup after every loose motion to a child

Unit_11_Communicable_disease_Gen_Nur.indd 251 09-03-2020 15:21:10