use of plenty of water, use of sanitary latrine, proper disposal of stools of young children and patients with infection, etc.

iii) Immunization : Immunization against measles is a potential intervention for diarrhea

Control Measures Two types of vaccines are used

a. Inactivated (Salk) vaccine (Injection)
b. oral (Sabin) polio vaccine (OPV).

iv) Fly control : Flies breeding in association with human or animal faeces should be controlled.

Treatment

1. Mild analgesics and sedatives to relieve pain and induce sleep.
2. For constipation – Mild laxatives
3. Antibiotics to prevent respiratory complication.
4. If respiratory failure occurs, treat with artificial respirators.
5. Rehabilitation

11.7 Diseases Transmitted Through Air

Chickenpox (Varicella)

Definition - Chickenpox or varicella is an acute highly infectious disease caused by varicella zooster

Causative organism

Varicella zoster

Mode of transmission - Droplet nuclei

Incubation period - About 10 -21 days

Complications (Children and Adults)

- Haemorrhages
- Pneumonia
- Encephalitis
- Acute cerebellar ataxia

The control measures are notifications isolation of cases for about 6 days after onset of rashes and disinfection of articles soiled by nose and throat discharges.

Clinical manifestations

Pre-eruptive stage:	Eruptive stage:
1. Fever	1. Rash, Clusters, itchy blisters
2. Pain in back	2. First appears on the trunk, then face, arms, axilla and in legs.
3. Shivering	3. Rashes in buccal surface.
4. Malaise	

Preventive measures

1. **Varicella zooster immunoglobulin** : Varicella zooster immunoglobulin given within 72 hours of exposure has been recommended for prevention of chicken pox.
2. **Vaccines** : The live attenuated varicella virus vaccine is safe and currently recommended for children between 12-18 months of age who have not had chicken pox.



Medication

- Anti – pyretic
- Anti viral drugs

Unit_11_Communicable_disease_Gen_Nur.indd 253 09-03-2020 15:21:11