1. Fever
2. Cold
3. Running nose
4. Irritating cough which gradually becomes paroxysmal within 1-2 weeks.

Complications

- Hernia
- Prolapsed rectum
- Sub conjunctiva hemorrhage
- Encephalopathy
- Pneumonia and bronchiectasis

Control measures

a. Early diagnosis
b. Isolation and treatment of cases
c. Disinfection of discharges from nose and throat are general principles of control
d. Early diagnosis is possible only by bacteriological examination of nose and throat Secretions.
e. Enthroning may help to shorten the duration of illness.

Treatment

Activation Immunization: It is a practice to administer pertussis vaccine in combination with diphtheria and tetanus toxoid as DPT vaccine.

It can be treated with antibiotics, usually erythromycin for 2 weeks.

Meaningococcal Meningitis

Definition

Meningococcal meningitis or cerebrospinal fever is an acute communicable disease caused by Neisseria. meningitis. It usually begins with intense headache,vomiting and stiff neck and progresses to coma within a few hours.

Causative organism

Neisseria meningitis

Mode of transmission - Droplet infection

Incubation period -2-10 days

Clinical manifestations

1. Headache
2. Fever
3. Nausea/vomiting
4. Photophobia
5. Neck stiffness and various neurological sign.

Complications

- Disseminated intravascular coagulation (DIC; blood clotting disorder)
- Encephalitis
- Persistent fever
- Seizures
- brain damage
- Behavioral and personality changes
- Vision loss (partial or total)
- Cerebral palsy
- Hearing loss (partial or total)
- Learning disabilities or mental retardation
- Paralysis (partial or total) Speech loss (partial or total)
- Severe bacterial meningitis also may cause the head and heels to bend backward and the body to bow forward (called opisthotonos),
- Coma, and death

Control Measures

a. Control of cases, carriers and contacts.
b. Cases: Treatment with antibiotics can save the lives of 95% of patients provided that, it is started during the first 2 days of illness. Penicillin is the drug of choice.
c. Contact: Close contact of persons with confirmed meningococcal disease patients

Unit_11_Communicable_disease_Gen_Nur.indd 258 09-03-2020 15:21:11