- Scarlatine form or petechial rash appears on third day of illness
- Head ache
- Nausea, vomiting
- Coffee colour vomiting
- Abdominal pain
- Pharyngitis
- Cough and dyspepsia

Dengue shock syndrome

- In addition to signs and symptoms of the above clinical feature client may go for shock.
- Sudden collapse
- Cold and clammy extremities
- Weak thread pulse
- Circumoral cyanosis along with haemorrhagic manifestation
- Occasionally epistaxsis, haematemesis, malena or subarachnoid haemorrhage.

Control measures

- Mosquito control

a. Cover all water containers.

b. Change water in flower vases every week.

c. Clean the surrounding area of the house.

d. Use insecticide spray in the house to kill adult mosquitoes.

- **Vaccines :** So far there is no satisfactory vaccine and no immediate prospect of preventing the disease by immunization

Other measures

- Isolation under bed rest
- Wear full sleeves shirts and full pants.
- Use of mosquito repellent creams, liquids, coils, mats etc.
- Use of bed nets for sleeping infants and young children during day time to prevent mosquito bite.

Treatment

a. Bed rest is advisable during the acute febrile phase.

b. Antipyretics or sponging are required to keep the body temperature at 98.6 degree F

c. Oral fluid and electrolyte therapy is recommended for patients with excessive sweating,vomiting or diarrhea.

d. Analgesics or a mild sedative may be required for those with severe pain.

e. Home available fluids to be given to prevent dehydration,

f. Fluid replacement should be minimum volume that is sufficient to maintain effective circulation during the period of leakage.

- **Management of shock :** Immediate replacement of plasma loss with isotonic salt solution.

Malaria:

Definition

Malaria is a protozoal disease caused by infection with parasites of the Genus plasmodium.

Causative organism

Female anopheles mosquitoes

Mode of Transmission - Malaria is transmitted by the bite of infected female anopheles mosquito.

Incubation Period - In some strains the incubation period may be delayed for as long as 6-9 months

Intermittent fever has 3 stages.

1. Cold stage (1/4 to ½ hours) - Head ache, Shivering fever rising rapidly
2. Hot stage (1/2 to 5 hours) - Very hot feeling, severe headache, skin flushed, fever starts falling

Unit_11_Communicable_disease_Gen_Nur.indd 262 09-03-2020 15:21:12