Head-Toe-Assessment

Assessment conducted by _____

LOC

☐ Alert ☐ Drowsy ☐ Lethargic ☐ Stuporous ☐ Coma

Orientation

☐ Person _____

☐ Place _____

☐ Time _____

Situation _____

Vitals

☐ Temp _____ ☐ R _____

☐ BP _____ Pulse Ox _____

Head

☐ Hair

☐ PERLA_____mm

☐ Nose _____

☐ Ears _____

☐ Mouth _____

- Midline tongue _____
- Moist _____
- Lesions _____
- Dentition _____

Neck

☐ Carotid Pulse _____ ☐ JVD + ☐ Trachea midline

Chest

☐ Apical pulse _____ ☐ Muffled ☐ Arrhythmia

☐ Breath sounds. Anterior _____

Posterior _____ Lateral _____

☐ Chest symmetry _____

☐ Skin Turgor (Clavicle) _____

Abdomen

☐ Inspection

☐ Ausculation

- LUQ (active / hypter / absent)
- RUQ (active / hypter / absent)
- LLQ (active / hypter / absent)
- RLQ (active / hypter / absent)

☐ Palpation _____

Upper Extremities

☐ Radial pulses equal, +2

☐Other: _____

☐ Temp vs. trunk (warm/cool)

☐ Grip equal and strong _____

☐ Capillary refill <3 sec

Practical 1-8.indd 297 09-03-2020 15:33:09