	Using the syringe: Feeding the following 1) Clamp the tube. Insert the tip of the large syringe with plunger, or bulb removed into the gastric tube. 2) Pour feeding into the syringe	
	3) Raise the syringe 12 to 18 inches above the stomach. Open the clamp. 4) Allow feeding to flow slowly into the stomach. Raise and lower the syringe to control the rate of flow. 5)Add additional formula to the syringe as it empties until feeding is complete	Gravity promotes movement of feeding into the stomach Controlling administration and flow rate of feeding prevents air from entering the stomach and nausea and abdominal cramping from developing
8	Termination of feeding: 1) Terminate feeding when completed. 2) Instil prescribed amount of water 3) Keep the client's head elevated for 20-30minutes.	To maintain patency of the tube Elevated position discourages aspiration of feeding solution into the lung
9	Mouth care: 1) Provide mouth care by brushing teeth 2)Offer mouthwash 3)Keep the lips moist	Mouth care promotes oral hygiene and provide comfort
10	Clean and replace equipment to proper place	To prevent contamination of equipment and prepare for the next procedure
11	Remove gloves and perform hand hygiene	To prevent the spread of infection
12	Document date, time, amount of residual, amount of feeding, and client's reaction to feeding. Sign the chart	Documentation provides continuity of care Giving signature maintains professional accountability

CONTRAINDICATIONS

Absolute contraindications

- Severe mid face trauma
- Recent nasal surgery

Relative contraindications

- Coagulation abnormality
- Oesophageal varices or stricture
- Recent banding or cautery of oesophageal varices
- Alkaline ingestion

REFERENCE

SanjitaKhadka et al,(2008) Fundamental of Nursing Procedure Manual, 1st edn. Japan International CooperationAgency (JICA): Nepal, pg - 104

Practical 1-8.indd 301 09-03-2020 15:33:09