In case of a child more than two years of age



1. Place the Electronic floor scale on the firm surface
2. Provide privacy
3. Undress the child allowing only minimal clothing (inner clothing).
4. Remove shoes/slippers
5. Adjust the scale to read zero
6. Request child or adolescent stand in the center platform of the scale
7. Ask the child remain motionless until the measurement can be obtained.
8. Measure the weight to the nearest 0.1 kg.
9. Check and compare previous weight
10. If the weight is in pounds and it must be converted to kilograms using conversion table.

Measurement of Length/Height

A child's length is measured in lying down position (recumbent). Height is measured in standing upright position.

Nursing Alert

If a child is less than 2 years old, measure recumbent length.

If the child is aged 2 years or older and able to stand, measure standing height.


In general, standing height is about 0.7 cm less than recumbent length.

If a child less than 2 years old will not lie down for measurement of length, measure standing height and add 0.7 cm to convert it to length. If a child aged 2 years or older cannot stand, measure recumbent length and subtract 0.7 cm to convert it to height.

Equipments needed

- Length board or Infantometer
- Height scale or Stadiometer

Measurement of recumbent length



1. Place a length board on a flat, stable surface such as a table.
2. Cover the length board with a thin cloth or soft paper for hygiene and for the baby's comfort.
3. Lay the child on his back with his head against the fixed headboard, compressing the hair.
4. Quickly position the head so that the child's eyes should be looking straight up.
5. Ask the mother or helper to move behind the headboard and hold the head in this position.
6. Stand on the side of the length board where you can see the measuring tape
7. Check that the child lies straight along the board and does not change position.
8. Shoulders should touch the board, and the spine should not be arched.
9. Hold down the child's legs with one hand and move the footboard with the other.
10. Apply gentle pressure to the knees to straighten the legs as far as they can go without causing injury.
11. If a child is extremely agitated and both legs cannot be held in position, measure with one leg in position.

Practical 1-8.indd 311 09-03-2020 15:33:10