Foot Care: The feet of ill bed ridden clients are easily susceptible to infection and other problems because the feet are far away from the heart than any other body part, they are most compromised by vascular conditions that interfere with normal circulation.

Conditions that generally affect bed ridden client include

1. Foot drop is a deformity in which the foot is extended abnormally at the ankle in the direction of the sole of the foot.
2. Intermittent claudications is a severe pain in the calf muscles caused by inadequate circulation. It usually occurs during walking, but subsides with rest.
3. Ulcers and gangrene are common side effects of diabetes. They occur because of inadequate circulation to the foot which retards natural healing process.

Oral Care: Cleansing of the mouth , teeth and gums is important to maintain the client's sense of well being as well as to prevent tooth decay and infection. Dental caries are the areas of localized destruction of tooth tissue by bacterial action. Caries are actually caused by acid production by bacteria which forms colony on the tooth surface.

f) Monitoring and surveillance:

The aspects of home health care which comes under monitoring and surveillance are

a) Neurological signs evaluation.
b) Urine glucose testing.
c) Vital signs.

Neurological Signs Evaluation: Neurological evaluation of the client can be obtained by objective and subjective data that are gathered through series of tests and evaluation techniques. The neurological status evaluation may be indicative of deteriorating condition or assessment of cognitive state. This is particularly important in home when traumatic injury is evaluated on when progressive neurological involvement may be side effect of medication therapy.

Urine glucose testing: Urine glucose testing is used to assess the status of person's diabetic condition. Diabetic results from body's inability to utilize food efficiently. When food is digested, it is broken into glucose, which is stored in liver and muscle tissue in the form of glycogen. Insulin facilitates the storage process. Diabetics do not produce sufficient insulin: therefore blood glucose levels rise to abnormally high levels. The normal fasting level of blood glucose is approximately 60 mg/dl to 115 mg/dl. Glucose does not appear in urine until the blood level reaches 180 mg/dl. Therefore, urine glucose level may be interpreted as reflection of actual blood glucose level.

Vital signs: Measurement of vital signs is done. To assess the physiological status of the client in relation to those vital canters of the body those are necessary to sustain life. The vital sign indicators are temperature, pulse, respiration and blood pressure. The temperature may be taken by oral, rectal or axillary route. The pulse may be measured by palpation, on auscultation of chest area; blood pressure is measured by means of sphygmomanometer.

Hot and cold applications: Hot and cold applications are applied to the clients in order to change the tissue temperature locally on systematically for a therapeutic purpose. Insulin injection: Since the major diabetes is thought to be the lack of inadequate use of insulin diabetic therapy often includes the use of insulin, in addition to dietary and exercise control. If the diabetic has little or no insulin production, capability in the pancreas, insulin is administered. The client family will be primarily responsible for performing the procedure on a daily basis. The injection should be performed with aseptic technique.

Intravenous therapy: Because of recent changes in health care industry encouraging early discharge from hospital, increasing

Practical 1-8.indd 326 09-03-2020 15:33:11