- Blockages in the blood vessels in arms or legs
- Diabetes
- High blood pressure
- Heart failure
- Kidney diseases
- Liver disease

Signs and Symptoms of ARF

- Oliguria - Urine output less than 400ml/day
- Anuria - Urine output less than 100ml/day
- Increased BUN creatinine
- Hyper Kalemia
- Metabolic acidosis
- Edema
- Mental changes
- Heart failure
- Nausea, vomiting
- Pruritus

Diagnosis

- History of underlying cause
- Blood- Increased potassium, BUN, Creatinine
- Urine: decreased volume and specific gravity is fixed or increased
- Renal ultrasound, renal scan, renal biopsy, CT scan or MRI, KUB X-ray, Retrograde pyelogram.

Management

- Removal of the underlying cause
- Diuretics
- Fluid restriction (600 ml plus previous day output) Calcium supplements or phosphate binding agents
- Nutritional therapy
  -Protein intake 0.6/kg/day
  -Potassium restriction
  -sodium restriction
  -Calorie intake 30 to 35 kcal/day/kg body weight
  -Dialysis

Chronic Renal Failure (ESRD-End stage Renal Disease)

Definition

Chronic or irreversible renal failure is a progressive reduction of functioning renal tissue or loss of renal function in which the body's ability to maintain metabolic and fluid electrolyte balance fails, resulting in uremia or azotemia over a period of months or years.

Causes

- Diabetic Nephropathy
- Hypertension
- Glomerulonephritis
- Pyelonephritis.
- HIV nephropathy
- Reflux nephropathy in children
- Polycystic kidney disease
- Kidney infections and obstructions
- Chronic pyelonephritis
- Head, Cadmium, mercury and chromium

Clinical manifestation

- Amenorrhea
- Testicular atrophy
- Malaise and Fatigue
- Pitting edema
- Periorbital edema
- Engorged neck veins
- CHF
- Anorexia
- Nausea
- Vomiting
- Seizure
- Constipation
- Peptic ulceration
- Diverticulosis

Unit_2_Medical_Surgery_Gen_Nur.indd 65 09-03-2020 13:13:01