Type 2 diabetes/Non-Insulin dependent diabetes mellitus(NIDDM)

Most common form of diabetes. Body produces insulin, but do not use it properly, glucose doesn't move into cells, they pile up in the bloodstream.

Risk factors



- Genetic
- Autoimmune
- Stress
- Environmental factors
- Obesity

Prediabetes

Slight elevation of blood glucose levels, regarded as indication that the person is at risk of progressing to Type 2 diabetes.

Gestational Diabetes Mellitus (GDM)

is defined as carbohydrate intolerance during pregnancy.

Risk factors

- Polycystic ovary syndrome
- women under age 25
- Hydraminos

Causes

- Family history of diabetes
- Overweight prior to pregnancy

Signs and Symptoms

- Hyperglycemia - Polyuria, Polydipsia, Polyphagia
- Weight loss
- Fatigue
- Blurred vision
- Poor wound healing
- Recurrent infection

Diagnosis

Urine analysis

- Glucose
- Ketone
- Microalbuminuria

Blood chemistry

- Blood glucose estimation, fasting and random blood sugar
- Oral Glucose tolerance test (OGTT)
- Check HbA1c (GLYCOSYLATED HAEMOGLOBIN LEVEL)

Management

Type 1

- Maintain and control sugar level
- Insulin therapy
- Healthy life style – exercise and diet.
- Islet transplantation
- Oral Antidiabetic agents
- Lipid control
- **DIET**
- Meal plan for Caloric restriction
- Weight reduction
- **EXERCISE:** Regularly scheduled, moderate exercise performed 30 to 60 minutes/day.
- Islet transplation

Type 2

- Maintain a healthy lifestyle
- Oral hypoglycemic agent and injection insulin if needed
- Dietary management and exercise

Gestational diabetes

- Insulin
- Physical activity
- Diet
- Plan Increase fiber intake

Unit_2_Medical_Surgery_Gen_Nur.indd 71 09-03-2020 13:13:02