Diagnosis

Diagnose uterine prolapse with

- A medical history and physical examination of the pelvis.
- Intra Venous Pyelogram (IVP)
- Renal sonography.
- X-rays.
- Ultrasound.
- Vaginal examination
- Rectal examination.



High post – pregnancy BMI raises pelvic organ prolapsed risk

Research suggests that having a high BMI after pregnancy increases the risk of a prolapse

Management

Prolapse up to the third degree may spontaneously resolve.

Complications

- Infertility
- Abortion
- Preterm labour
- Risk of operative delivery
- Anaemia due to heavy bleeding
- Uterine cancer

Vaginal pessary: It is a removable device placed into **the vagina**.

Surgery: Hysterectomy

Nursing management

Teach and insist to practice Kegel exercise during pregnancy and post natal period


Exercise

Mild uterine prolapse can be treated with Kegel exercises

- Tighten the pelvic floor muscles, as you are attempting to stop urinating and hold for 5 seconds
- Take a 5-second break and repeat for three to 10 times per day.
- Avoid more child birth.

Prevention

- Maintain optimal weight.
- Avoid constipation by eating a high-fiber diet.
- Perform Kegel exercises to strengthen the pelvic muscles.
- Avoid heavy lifting or straining. Preventing and treating constipation
- Avoid chronic cough



Unit_2_Medical_Surgery_Gen_Nur.indd 78 09-03-2020 13:13:03