







Table 2.1 Vegetable Dyes

Name of the Plant	Parts used	Dyed fabric
Indigo		
Indian madder		
Turmeric		
Marigold		
Henna		
Tea		
Onion		

DO YOU KNOW?

Who is Adolf Von Baeyer ?



Adolf Von Baeyer

Adolf Von Baeyer is a German chemist, who was awarded the Nobel Price in Chemistry in 1905 for discovering the molecular structure of Indigo and developing a process to produce it synthetically.

2.2.3 Animal Dyes

Dyes extracted from certain insects and invertebrates are called as animal dyes. Various shades of red and purple were obtained from animal origin. Cochineal, Tyrian purple and Lac are the commonly used animal dyes.

1. Cochineal

   Cochineal dye is extracted from the dried bodies of the female red bug (*Dactylopius coccus*). It produces crimson and scarlet colours with mordants aluminium and tin oxide. This dyestuff was mostly used for dyeing wool and silk. These dyes exhibit excellent fastness properties.

2. Tyrian Purple

   This dye is extracted from the sea snails found in Mediterranean Sea. The amount of dye produced was very limited and therefore very expensive. Hence, it is called Royal purple.

3. Lac Dye

   This dye is extracted from the fluid secreted by the lac insect (*Lauifer lacca*), which lives on the twigs of the banyan trees and other varieties.

XII TDD_English version CHAPTER 02.indd 22 1/13/2020 2:43:59 PM