Sources

Green leafy vegetables, cereals, fruits. Synthesized by bacteria in gut.



Functions

- Important component in blood coagulation
- Helps in formation of blood clotting factors



Newborn babies have sterile intestinal flora and cannot produce Vitamin K, hence given artificially.

Deficiency Syndrome: Alteration in blood clotting mechanism

II. Water Soluble Vitamins

Vitamin B_1- Thiamine	Vitamin B_2- Riboflavin	Vitamin B_3 - Niacin
Vitamin B_5- Pantothenic acid	Vitamin B_6- Pyridoxin	Vitamin B_7- Biotin
Vitamin B_9- Folic acid	Vitamin B_{12}- Cyanocobalamine	Vitamin C - Ascorbic acid

These vitamins are essential for many metabolic functions of the body. These are daily required in diet.

Vitamin B_1- Thiamine

Thiamine is essential for the normal metabolism of carbohydrates and fats. It is necessary for the transmission of nerve impulses.

Daily requirements: 0.5 to 2 mg

Sources

whole grain cereals, wheat, ragi, pulses (dhal), vegetables and potatoes, green leafy vegetables. Meat, fish, liver and eggs.





Polished milled rice and wheat flour of white variety are deficient of vitamin B_1.

Functions

- Helps in carbohydrate utilization.
- Maintenance of appetite and digestion.

Deficiency Disorder

Beriberi- condition in which there is a severe muscle wasting, growth retardation in children, neurological disturbances and frequent infection.



XII_Nursing_Vocational_Unit 4.indd 115 1/11/2020 2:40:15 PM