Foods to be restricted

- High fibre foods like whole grain cereals, whole wheat flour and pulses
- All raw vegetables and fruits except papaya and banana
- Fried fatty foods such as samosa, pakodas, halwa etc
- Chemical irritants such as condiments, spices, pickles, chutneys etc should be avoided

Diet in Constipation

Constipation is decreased frequency of passing stools or complete retention of faeces. The diet should include foods rich in fibre content like whole cereals, legumes and vegetables. Other foods such as fruits rich in fibre e.g. banana, apple, guava etc. more of fluids should be consumed in any form.

STUDENT'S ACTIVITY

A man complaints of constipation. What foods should you tell him to include in his diet?

Plenty of water

Drink more water and other liquids, such as natural fruit and vegetable juices and clear soups, to help the fibre work better. This makes your stools softer and easier to pass.

Drinking enough water and other liquids is also a good way to avoid dehydration. Staying hydrated is good for your overall health and can help you avoid getting constipated.

Diet in Diarrhoea

Diarrhoea is increased frequency of loose or watery stools. It occurs in infectious condition of colon. Diet must include mainly fluids like oral boiled water containing electrolyte salts, i.e glucose and sodium chloride. Fluids should be given frequently. Gradually fruit juices, butter milk, diluted milk must be introduced. During treatment phase soft bland diet like milk pudding, rice dhal gruel etc. can be started. Oral rehydration solution must be given which fulfills the nutritional and electrolyte balance.

Oral Rehydration Solution

Home Made ORS recipe - The "simple solution" Preparing.

One (1) Litre Oral Rehydration Solution [ORS] using Salt, Sugar and Water at Home.

Ingredients

Half (1/2) level teaspoon of Salt.


Six (6) level teaspoons of Sugar.

One (1) Litre of clean drinking or boiled water and then cooled 5 cupfuls (each cup about 200 ml).

Preparation Method: Stir the mixture till the salt and sugar dissolve.

6 Level Tea Spoons of Sugar

Half Level Tea Spoon of Salt



1 Litre of Water - 5 Cups
(each cup about 200 ml)

A child with diarrhoea should receive oral rehydration salts (ORS) solution and a daily zinc supplement for 10–14 days. Diarrhoea medicines are generally ineffective and can be harmful.

XII_Nursing_Vocational_Unit 4.indd 127 1/11/2020 2:40:38 PM