Diet in Renal Failure

Renal failure is accumulation of urea in the blood and tissues. It may be acute or chronic. In this condition glomerular function is affected. The diet should be low in protein content, sodium, potassium and high in calories. Salt should be restricted.

Foods to be included

- Milk and its products in needed quantity.
- Ghee, oil, sugar, potatoes and other starchy foods can be used to increase the palatability of foods.
- Fruits without potassium can be taken.
- Spices and condiments in small amounts can be taken.

Foods to be avoided

- Meat, poultry and fish.
- Dry fruits like almonds, peanuts, Cashewnuts and walnuts.
- Extra pulses, cereals, legumes, peas and beans.
- Cakes, biscuits and bakery products.
- Fruits if potassium is restricted
- Green leafy vegetables if potassium is restricted

STUDENT'S ACTIVITY

Discuss the various foods that can be given to
a) Diabetes mellitus
b) Constipation
c) Renal failure

4.10 METHODS OF COOKING

Cooking is an art. It is linked with the dietary and cultural pattern of people. Almost all foods consumed need some form of cooking and processing before they are fit for serving and consumption. Fruits and some vegetables used in salad or chutneys are consumed uncooked.

Cooking has many Benefits

1. Cooking increases palatability of the food
2. It makes mastication easier and renders the food easy to digest
3. It sterilizes food by killing microorganism
4. Adds new flavor and stimulates digestive juices
5. Good cooking increases the acceptability of food
6. It improves the appearance of food

Methods of Cooking

- Boiling
- Simmering
- Steaming
- Stewing
- Roasting
- Frying
- Baking
- Grilling
- Solar cooking

Boiling: Cooking in water at its boiling point 100°c is known as boiling. Most of the vegetables are cooked by this method. Boiling in excess water may result in loss of vitamins and minerals. Specific time and minimum water must be kept for boiling these items.



XII_Nursing_Vocational_Unit 4.indd 129 1/11/2020 2:40:39 PM