


2. **Lateral Palpation / Lateral Grip** (both Right and left lateral) - Helps to locate fetal back and limbs



3. **Pelvic Grip I / Superficial Pelvic Grip** -Helps to determine whether head or breech is presenting at pelvic brim. Whether the presenting part is engaged / fixed / free.



4. **Pelvic Grip II / Deep Pelvic Grip** - Helps to know the degree of flexion of head.



Check or ask for Foetal Movements

- Fetal movement are reliable sign of foetal well - being.
- These are felt around 18-22 wks of pregnancy (felt earlier in multigravida than primigravida).
- Normally 10-12 foetal movements should be felt by the pregnant woman in a day.
- Decreased foetal movements may be an indication of foetal distress.
- Pattern of foetal movement may change prior to labour due to reduced space.
- If Foetal Movements are absent or not felt, consult ANM or doctor.

3. Auscultation

- Use a fetoscope or stethoscope.
- Best heard on the side of the back of the fetus.
- In vertex presentation FHS is best heard midway between the line joining the umbilicus and the anterior superior iliac spine on the side of the back.
- In breech presentation FHS is heard above the umbilicus (Fetal Heart Sound).
- Count the FHS for one full minute FHR (Fetal Heart Rate).



- FHS is heard over the abdomen by stethoscope / fetoscope after 24 weeks of pregnancy.
- Normal FHR is 120 – 160 beats per min.

XII_Nursing_Vocational_Unit 5.indd 162 1/11/2020 2:46:17 PM