- FHR < 120 beats per min or > 160 beats per minute, indicates fetal distress.
- Confirm that you are listening to the FHS and not maternal pulse.

DIET in Pregnancy

- Total caloric intake should be increased to 300 kcal /day due to 15% increase in BMR.
- Diet should contain 20% Protein (better from animal source), 30% fat, and 50% carbohydrates.
- Sufficient fluids should be taken. (10 glasses for a day).
- Absorption of iron is interfered if taken with tea, coffee or foods rich in fluoride.
- Enhanced if taken with lemon water or orange juice.
- Encourage mother to take plenty of fruits and vegetables like mango, guava, orange, amla etc containing vitamin C.
- Emphasize the importance of high protein diet like black gram, ground nuts, whole grains, milk, eggs etc.

Supplementation - **Iron, folic acid and calcium as prescribed**

WEIGHT gain in Pregnancy

- Total weight gain approximately 12 Kgs.
- Weight gain of 2 kgs in first trimester. 5 kgs in second and 5 kgs in third trimester.
- Monitoring of weight gain should be done in conjunction with close monitoring of BP.
- Overweight or sudden increase of weight is to be notified immediately.

Oral Care

- It is easy to have an increase in dental decay cavities due to pregnancy. Heartburn, increased snacking, morning sickness can all increase chances of developing tooth decay or gum disease. Good oral care is an important during pregnancy as it is the most important time of life.

CARE OF BREAST

- Breast engorgement may cause discomfort during late pregnancy. A well-fitting brassiere can give relief.
- Travelling during pregnancy is not prohibited but some precautions must be taken.
- Avoid long trips if possible. Always check with care provider before travelling. Wear seatbelt, the shoulder belt shouldd go between the breasts and the lap belt should go under the tummy. Plan for frequent stops. Get out and walk as much as possible. This will prevent swelling of the foot.
- When travelling by air, need to drink extra fluids. Walk around whenever possible. Do isometric exercises of legs and foot to help prevent swelling and blood clots. Do not plan to travel after 34 weeks of pregnancy.

5.10 ANTENATAL EXERCISES

Uses of Antenatal Exercise

1. Good muscle tone is maintained during labour and postnatal period.
2. Helps to reduce the backache.
3. Assist in effective pushing during labour.

Types of Antenatal Exercise

- Deep breathing exercise.



STUDENT'S ACTIVITY

Ask the students to demonstrate the Antenatal exercises to each other.

XII_Nursing_Vocational_Unit 5.indd 163 1/11/2020 2:46:18 PM