Medications during Pregnancy

- ❖ Avoid over the counter medications
- ❖ Inj. T.T. (Tetanus Toxoid) For preventing maternal and neonatal tetanus
  - Dosage schedule: Two doses 0.5 ml I/M in upper arm
  - First dose - As soon as woman registers for ANC
  - Second dose - 4 weeks after the first dose even if it is after delivery
  - Inform that there may be slight swelling, stiffness or pain at the injection site
- ❖ Iron and folic acid supplementation
  - 1 tablet / of IFA (100 mg elemental iron+0.5 mg of folic acid)
  - Duration: At least 100 days during pregnancy
  - Continue for 3-6 months in postpartum period
  - Folic acid supplementation (400 µg) upto 12 weeks of pregnancy to prevent Neural tube defects
- ❖ Regular consumption of iodized salt
- ❖ Calcium supplementation (500mg-1g / day)



- Iron Tablets to be taken regularly preferably early morning on an empty stomach
- If nausea or pain abdomen, it may be taken after meals or at night
- Stools will be dark in colour while on tablets
- May have constipation, relieved by taking plenty of fluids and roughage

Hazards during Pregnancy

Hazards affect the normal functions of the mother and unborn baby's body.

They can cause birth defects in the unborn child. Hazards include physical factors, germs, tobacco, alcohol, health conditions of the mother, chemicals, and drugs.

Warning signs of Pregnancy

- ❖ Vaginal bleeding
- ❖ Persistent vomiting
- ❖ Chills and fever
- ❖ Sudden escape of clear fluid from vagina
- ❖ Abdominal or chest pain
- ❖ Increase or decrease in fetal movement

The nurse must educate the women that if above symptoms are present they must come to the hospital immediately.



STUDENT'S ACTIVITY

Prepare health education plan for Primi Antenatal mother

Minor disorders in pregnancy

- ❖ Morning sickness
- ❖ Heartburns
- ❖ Varicose veins
- ❖ Backache
- ❖ Breathlessness
- ❖ Palpitations
- ❖ Vaginal discharge
- ❖ Constipation



- Nearly 4 million newborns die {40% of under 5 deaths} within 28 days of birth
- Three quarters of neonatal deaths occur during first 7 days
- For every newborn death, 20% from birth injury, complications of preterm birth or other neonatal conditions
- A child born in a least developed country is 14 times more likely to die within first 28 days of life as compared to industrialized country

XII_Nursing_Vocational_Unit 5.indd 165 1/11/2020 2:46:18 PM