- Pertussis
- Tetanus
- Measles

b) Other vaccine Preventable Diseases

- Hepatitis B
- Rubella
- Typhoid
- Influenza
- Chickenpox
- Yellow fever
- Plague
- Rabies
- Mumps
- Haemophilus influenza Type B
- Meningococcal meningitis
- Pneumococcal Pneumonia
- Rotta virus
- Diarrhoea
- Cholera
- Malaria

6.7.3 The Cold Chain

The cold chain system is necsessary, because vaccine failure may occur due to store and transport under strict temperature controls.

6.7.2 Immunization Schedule

National Immunization Schedule for infants, Children and pregnant Women				
Vaccine	**When to give**	**Dose**	**Route**	**Site**
For Pregnant Women				
TT – 1	Early in pregnancy	0.5 ml	Intra-muscular	Upper Arm
TT – 2	4 weeks after TT – 1	0.5 ml	Intra-muscular	Upper Arm
TT - Booster	If received 2 TT doses in a pregnancy within last 3 yrs	0.5 ml	Intra-muscular	Upper Arm
For Infants				
BCG	At birth or as early as possible till one year of age	0.1ml (0.05 ml till 1 month age)	Intradermal	Left Upper Arm
Hepatitis B	At birth or as early as possible within the first 24 days	0.5ml	Intra–muscular	Antero–lateral side of mid-thigh
OPV – O	At birth or as early as possible within the first 15 days	2 drops	Oral	Oral
OPV 1,2&3	At 6 weeks 10 weeks & 14 weeks	2 drops	Oral	Oral
DPT 1,2&3	At 6 weeks 10 weeks & 14 weeks	0.5ml	Intra-muscular	Antero–lateral side of mid-thigh
Hep B 1,2&3	At 6 weeks 10 weeks & 14 weeks	0.5ml	Intra-muscular	Antero–lateral side of mid-thigh
Measles	9 completed months – 12 months	0.5ml	Subcutaneous	Right Upper Arm
Vitamin – A (1st dose)	At 9 months with measles	1ml (1 lakh IU)	Oral	Oral
For Children				
DPT booster	16-24 months	0.5 ml	Intra-muscular	Antero–lateral side of mid-thigh
Measles 2nd dose	16-24 months	0.5 ml	Subcutaneous	Right Upper Arm
OPV Booster	16-24 months	2 drops	Oral	Oral
Japanese Encephalitis	16-24 months	0.5 ml	Subcutaneous	Left Upper Arm
Vitamin – A (2nd to 9th dose)	16 months. Then, one dose every 6 months upto the age of 6 years	2ml (2 lakh IU)	Oral	Oral
DPT Booster	5-6 years	0.5 ml	Intra-muscular	Upper Arm
TT	10 years & 16 years	0.5 ml	Intra-muscular	Upper Arm

JE vaccine, in selected endemic districts after the campaign.

The 2nd and 9th dose of Vitamin A can be administered to children 1 to 5 years old during bi-annual rounds, in collaboration with ICDS.

XII_Nursing_Vocational_Unit 6.indd 196 1/11/2020 2:50:47 PM