Common problems	Management
Dehydration Dehydration results when there is more loss of fluid in the body than the intake. This happens in severe diarrhoea and vomiting,which is more serious in children. **Signs of dehydration:** • Little or no urine • Sudden weight loss • Dry mouth • Sunken, tearless eyes • Loss of elasticity or stretching of the skin • Very severe dehydration may cause rapid, weak fast pulse, deep breathing, fever, or fits	**Prevention and treatment of dehydration:** • The person should drink large amount of liquids, water, tea, soup etc., • The person should be given plenty of liquids or rehydration to drink first • A mother can prepare rehydration drink • A 4-finger scoop of sugar or jaggery and 3 pinch of salt and add it to one litre of boiled and cooled water • Give the person sips of drink every five minutes, day and night, until he begins to urinate normally. A small child needs atleast one litre a day
Contents of rehydration packet Marketed as ORS under different trade names Glucose-20.0gm • Sodium chloride - 3.5 gm • Sodium bicarbonate - 2.5 gm • Potassium chloride - 1.5 gm 	To prepare oral rehydration salt solution: • Wash hands • Pour one litre of clean water into clean container • Open the sealed packet and pour the ingredients and mix well • Fresh ORS should be prepared each day. Container should be kept covered
Contents of sugar salt solution; Sugar - 40gm or honey/glucose 20 gm Common salt - 5 gm (one pinch) One litre of water - after mixing well in cool drinking water, the solution should have the taste of tears.	• Keep giving rehydration drink often in small sips, even if the person vomits • If the person vomits all the drinks shift the person to the health centre to give fluids through veins.

XII_Nursing_Vocational_Unit 1.indd 11 1/11/2020 1:59:13 PM