to diarrhoea. All the deaths due to diarrhoea can be prevented by treating dehydration, with the use of ORS (Oral Rehydration Solution) and administration of zinc tablets along with adequate nutritional intake by the child. ASHA worker would undertake the distribution of ORS packets to households with the supervision of safe drinking water, breast feeding, sanitation, nutrition and hand washing. (ASHA – Accredited Social Health Activist)

6.10.8 The Reproductive and Child health Programme (RCH)

This programme was launched in October 1997. The main aim of the programme is to reduce infant, child and maternal mortality rates and to reduce total fertility rate.

The package include

- Prevention and management of unwanted pregnancy.
- Provision of services to promote child survival.
- Prevention and treatment of reproductive tractt infection and diseases.

WHO on 24th February 2012 removed India from the list of countries with active endemic wild polio virus transmission.

6.10.9 Integrated Child Development Services (ICDS)

It was initiated by the Government of India in the Ministry of social and women's welfare in 1975. Which provides food, preschool education and primary health care to children under 6 years of age and their mothers.

6.9.10 Balwadi Nutrition Programme

This programme was started in the year 1970 under the Ministry of Social Welfare. The aim is to provide primary education to 3-6 years children. The food supplement provides 300Kcal and 10gms of protein per children.

6.10.11 Mid-day meal programme

Mid-day meal programme is also known school lunch programme to noon meal programme.

This Programme has been in operation since 1961 throughout the country under Ministry of Education. The object is to attract more children for admission to schools and return them so that literacy improvements of children could be brought about.

Principles

- The meal should be a supplement and not a substitute to the home diet.
- The cost of the meal should be reasonably low.
- The menu should be frequently changed to avoid monotonous.

6.10.12 Pallisirar Kannoli Kappom Thittam:



Refractive errors (need for glasses) are the second main cause for avoidable blind and the rate of optometrists in reducing this condition

XII_Nursing_Vocational_Unit 6.indd 203 1/11/2020 2:50:51 PM