Common problems	Management
Vomiting • Many person, especially children have an occasional stomach upset with vomiting. Often no cause can be found • There may be mild stomach or guts, such as infection, poisoning from spoiled food • Any sickness with acute fever or severe pain may cause vomiting, especially malaria, hepatitis, tonsillitis, earache, meningitis, urinary infection, gallbladder pain or migraine headache	To help control simple vomiting: • Sip some tea with sugar but without any milk. Adding ginger or lime juice may also help • For dehydration give small frequent sips of cool tea, or rehydration drink • If vomiting does not stop soon, use vomit control medicines like promethazine, diphenyldramine or phenobarbitol
Diarrhoea and dysentery • When a person passes loose or watery stools more than two or three times it is called diarrhoea. If mucus and blood can be seen in the stool it is dysentery • Diarrhoea can be mild or serious. It can be acute or chronic • Diarrhoea is more common and more dangerous in young children **Causes of diarrhoea:** • It has many causes • Poor nutrition • Virus infection or intestinal flu • An infection of the gut caused by bacteria, amoeba, urinary infections • Malaria • Food poisoning • Allergic to certain foods • Side effects produced by certain medicines, such as ampicillin or tetracycline	**Prevention of diarrhoea:** The prevention of diarrhoea depends on good nutrition and cleanliness. • Exclusive Breast feeding rather than bottle feeding for the first six months • When you begin to give the baby new or solid foods • Keep the baby and the surrounding clean **Treatment of diarrhoea:** For most of the cases of diarrhoea no medicine is needed. If the diarrhoea lasts along times, the bigger danger is malnutrition. so the most important part of treatment has to do with giving enough liquids and good food, no matter. **Prevent or control dehydration:** A person with watery diarrhoea must drink large amount of liquids. • Give him rehydration drink • *Meet nutritional need* A person with diarrhoea needs food. This is especially important in small children or persons who are poorly nourished.

XII_Nursing_Vocational_Unit 1.indd 12 1/11/2020 1:59:13 PM