
Diagnosis

- History collection
- Clinical examination including mental function
- CSF examination
- CT scan

Treatment

- Home health aid to assist in personal hygiene home cares to assist in house work transportation
- Support groups such as the alzhemeirs associations are often of value to the family
- Legal council should be recommended to help the patient and family

7.5.3 Depression



Depression occurs in 5 to 10% of community - dwelling elderly due to lack of social supports.

Clinical Manifestations

- Sleep disturbance
- Lack of interest
- Feeling of guilty
- Suicidal ideation

Treatment

- Medical and pharmacological Management Methylphenidate 5 to 10 mg at 8 a.m. and noon is often very effective.
- Treatment should be continued for up to a year because relapse rates are higher in the elderly.

STUDENT'S ACTIVITY

Identify the physical and psychological changes of your grandmother who is above 75 years. Based on the observation how wil you provide a care

7.5.4 Acute Confusional State (Delirium)

- Delirium called acute confusional state and progressive to disorientation.
- Delirium due to hyponatraemia, hyperosmolality, hypercapria



Clinical Manifestations

- Disturbance in memory,
- Thought disorder
- Emotional changes
- Disturbance of perception.

XII_Nursing_Vocational_Unit 7.indd 213 1/11/2020 3:01:45 PM