TREATMENT

Supportive treatment ECT (Electroconvulsive Therapy)

Anopsychotics

7.6 SENSORY SYSTEM

7.6.1 Sensory Problems

- Decreased visual perception
- Decreased elasticity of the ear drum
- Decreased sense of smell
- Decreased taste perception
- Decreased touch sensation

People interact with the world through the sense, sensory losses associated with old age affect all sensory organs. Nearly half of older man and one third of older women report difficulty hearing without aHearing Aid 16% of old men and 19% of older women report difficulty seeing even with the corrective lenses.

7.6.2 Vision



- Decrease peripheral vision.
- Decrease night vision.
- Decrease capacity to distinguish colour.
- Reduced lubrication resulting in dry, itchy eyes.

Management

- Approach people directly not from the side.
- Proper lighting can make a big difference.
- Assist with choosing clothes if needed
- Use of natural tear products.

7.6.3 Hearing



- Auditory changes begin to be noticed at about 40 years of age.
- Environmental factors such as exposure to noise, medications and infections as well as genetics may contribute to hearing loss.
- Hearing loss may cause older people to respond inappropriately.
- Older people often cannot following conversation because tones of high frequency consonants (F, S, TH, CH, SH, B, T, P) all sound alike.

Management

- Reduce extra noise when trying to have a conversation.

XII_Nursing_Vocational_Unit 7.indd 214 1/11/2020 3:01:48 PM