Priority	Colour of the tag	Condition
First (Immediate)	Red	Life threatened. have a high probability of survival
Second (Intermediate)	Yellow	Seriously injured Can delay treatment and transport for 2 hours
Third (waiting)	Green	Walking wounded Can delay transport 3 hours
Low priority	Black	Expectant, died victims, severely injured 100%

husbandry up-stream of intake points in rivers and streams

- Upgrade wells to ensure that they are protected from contamination
- Estimate the maximum yield of wells and if necessary ration the water supply
- All water tankers should be cleaned and disinfected before transporting water

Basic Sanitation and personnel hygiene

Many communicable diseases are spread through faecal contamination of drinking water and food. Hence every effort should be made to ensure the sanitary disposal of excreta. Emergency latrines should be made available to the displaced, where toilet facility have been destroyed, washing, cleaning and bathing facility should be provided to the displaced persons.

Vector Control

Vector borne diseases should be intensified in the emergency and rehabilitation period, especially in areas where such disease are known to be endemic. Special concerns are Dengue fever and Malaria, Leptospirosis and rat bite fever, typhoid and plaque. Flood water provides chance for breeding opportunities for mosquitos.

8.3.4 Role in disaster preparedness



- Facilitate preparedness
- Initiate and update disaster plan
- Provide educational programme in specific area
- Organize disaster drills (mass drills)
- Provide updated record
- Educate the vulnerable population
- Nurse should seek safe environment
- Assess the environmental hazard
- Understand the community resources
- Physical readiness
- Professional readiness
- Community readiness.

8.4 CORE EMERGENCY PREPAREDNESS

The term emergency refers to any extraordinary event or situation that requires

XII_Nursing_Vocational_Unit 8.indd 234 1/11/2020 3:03:19 PM