

TAB. PARACETAMOL is dangerous to take more than eight 500 mg or six 650 mg tablets in 24 hours as you may damage your liver, which may be irreversible and fatal.

Drug dependence: Drugs capable of altering mood and feelings on repeated use of such drugs. E.g., morphine

Iatrogenic disease: This a physician induced or drug induced disease. E.g., peptic ulcer by salicylates.

Drug allergy: It is an immunologically mediated unpredictable reactions which are not related to the therapeutic effects.

Carcinogenicity and Mutagenicity: It is the capacity of a drug to cause cancer. e.g., anabolic steroids.

9.10 STORING OF MEDICINE AND SAFETY MEASURES

1. Most medications should be stored in a dry place away from heat and humidity.
2. If the medication needs to be protected from light, it should be stored in a container that keeps out light.
3. If the medication needs to be refrigerated, and you do not have a fridge available, use a cooler with an ice pack.
4. Keep track of the expiry dates of the medications by asking the pharmacy team to put this information on your medication vial.
5. Don't store multiple medications in the same vial, as this makes it hard to keep tracing of which medications are which, which doses have been taken, and what are the expiry dates.
6. Put all medicine up and away, out of children's reach and sight.
7. Close your medicine caps tightly after every use.
8. Choose child-resistant caps for medicine bottles. If pill boxes or non-child resistant caps are the only option, it's even more important to store these containers up high and out of sight when caring for kids.

9.10.1 How to administer medicine Safely

1. Use the dosing device that comes with the medicine. Proper dosing is important, particularly for young children.
2. Kitchen spoons aren't all the same, and a teaspoon or tablespoon used for cooking won't measure the same amount as the dosing device.
3. Keep all medicine in their original packages and containers.
4. Take the time to read the label and follow the directions. Even if you have used the medicine before, sometimes the directions change about how much medicine to give.



The important life saving drug also has its own side effects.



XII_Nursing_Vocational_Unit 9.indd 249 1/11/2020 4:11:47 PM