- Kidney tray and paper bag
- Adhesive tapes

Procedure:

1. Wash hands.
2. Measure the length of the catheter from the tip of the nose to ear lobe. Mark the length with ink.
3. Check the cylinder for the working condition. Open the main valve in an anti clock wise direction. Look for the pressure reading on the gauge. Adjust the flow of oxygen to 2-4 Litres/min for adults.



4. When the wheel valve is opened the oxygen will start bubbling through the water in the humidifier.
5. Lubricate the tip of the catheter sparingly with water soluble jelly and check the flow by immersing it in water.
6. Introduce catheter slowly into one of the nostrils of the previously marked distance. Never use force.
7. Check the position of the catheter in the oropharynx at the level of the uvula.
8. It can be checked by asking the patient to open his mouth widely.
9. Depressing the tongue with tongue depressor and use the flash light to see the throat
10. Fix the catheter over the forehead or at the cheek of the patient with adhesive tapes
11. Save the connecting tube to the bed clothes or patient gown with safety pin

After care of the patient and articles:

- Stay with the patient till he is at ease
- Keep the patient warmth and comfortable
- Assess the vital signs frequently
- Record the procedure with date and time in the nurses record
- Check the cylinder for its good working condition
- Change the nasal catheter by every 8 hours
- When the oxygen is to be stopped, do it gradually
- Watch the patient for any deteriorating symptoms after the removal of oxygen



XII_Nursing_Vocational_Practical.indd 301 1/11/2020 4:29:33 PM