Foods to be excluded from diet:

- Sugar, honey, jaggery, sweet syrup and fried foods
- All roots
- Cool drinks
- Horlicks, Bournvita etc
- Coconut, coconut oil, palm oil
- Ground nut and all nuts
- Tinned food
- Alcohol
- Butter, ghee, cheese

Snacks to be eaten in between meals:

- Butter milk
- Vegetable soup
- Vegetable salad
- Lime/tomato juice without sugar

Hypertension: (High blood pressure)

This is a common condition in middle age. Especially in obese people. It can lead to complications such as heart disease, kidney disease and stroke.

To prevent and treat hypertension:

1. Over – weight people should lose weight
2. Low – fat, salt – free diet is needed.
3. Coffee and other stimulants should be avoided.
4. Tell the person to relax and avoid tension.

A dietary menu for hypertensive patients is as follows:

Morning	Weak Tea one Cup
Break fast	Bread or idli with sugar, or jam fruit one cup, skimmed milk one cup. Nuts one table spoon.
Mid morning	Fruit juice 200 ml
Lunch	Rice or chappati / one serving dhal or meat or fish, one cup vegetable, one serving skimmed butter milk or pudding one cup
Evening	Biscuits two and fruit juice one glass and roasted nuts two table spoons
Dinner	Similar to lunch

Diet

Definition: Diet is an important as medicine in the treatment of diseases. A modification in the diet or in the nutrients can cure certain diseases. E.g a patient suffering from peptic ulcer needs a bland diet for this recovery. A salt free diet can reduce the blood pressure in a patient with hypertension.

Types of Diet:

- Full diet
- Liquid diet
- Soft diet
- Semi solid diet
- Bland diet
- High protein diet
- Low protein diet
- Salt free diet or salt low diet

XII_Nursing_Vocational_Practical.indd 318 1/11/2020 4:29:44 PM