
4. In severe diarrhoea Antibiotics [**Cefixime**, **ceftriaxone**, and **cefotaxime**] resolve the symptoms of diarrhoea

Preventive measures



2.2.3 Typhoid



It is an acute illness, associated with constant high fever, which is caused by bacteria Salmonella typhi. This is transmitted through water and food and found in the feces of the infected person. Other name of typhoid fever is entric fever .

Causative organism

- Salmonella typhi

Incubation period

- 8 to 14 days

Symptoms of Typhoid

- Loss of appetite
- Weakness
- Abdominal pain
- Vomiting
- Diarrhoea or constipation
- High fever
- Head ache

Management


Antibiotics [Trimethoprim-sulfamethoxazole, Ampicillin and Ciprofloxacin]. are the most important to treat the infection, for the complete removal of the bacteria.

Typhoid vaccines are recommended for where there is a risk of typhoid is very high.

Preventive Measures

- Drink pure and safe water
- Avoid raw fruits and vegetables
- Street food to be avoided
- Wash hand before and after using toilets.

2.2.4 Hepatitis A



It's the most highly contagious liver infection, which is caused by hepatitis **A** virus. This virus cause inflammation of liver and affects the ability of liver function.

Causative organism

- Hepatitis A virus[HAV]
- Hepatovirus

XII_Nursing_Vocational_Unit 2.indd 28 1/11/2020 2:25:08 PM