- Airborne
- Gastrointestinal infection

Signs and symptoms

- Cutaneous infection
- Malaise
- Fever
- Headache
- Respiratory infection
- Cyanosis
- Shock
- Death
- Nausea
- Loss of Appetite
- Vomiting
- Abdominal pain
- Toxemia

Risk factors

- Workers who come into contact with animal hides
- Workers involved in the handling of dead animals

Diagnosis

- Identification of the typical Gram positive bacilli of Bacillus anthracis from skin lesions
- Respiratory secretions

Management

- Treatment should be initiated as soon as possible.
- Mild cases of cutaneous anthrax may be effectively treated with oral Penicillin or tetracycline
- Treatment of inhalation or gastrointestinal anthrax requires high dose intravenous Penicillin

Preventive Measures

- Controlling anthrax in livestock
- Preventing gastrointestinal anthrax by forbidding the sale of meat from sick animals or animals that have died of the disease
- Regular cleaning and disinfection of the workplace
- Disinfecting animal products
- Processing hides, wool, and bone by tanning, dyeing, carbonizing or using acid treatment
- Administering vaccines

Good hygiene practice (hand washing avoiding hand to eye/mouth contact) and need to cover cuts and abrasions.

2.6.2 Tetanus

Tetanus is caused by the infection of wounds by bacterium Clostridium tetani, which produces toxin that damages the nervous system and muscles



Causative organism:
Bacterium Clostridium Tetani

Incubation period

3–21 days

Signs and symptoms

- Muscle rigidity

XII_Nursing_Vocational_Unit 2.indd 41 1/11/2020 2:25:26 PM