## Symptoms

- Dry , flaky skin (xeroderma)
- Chronic fatigue
- Fever that comes and goes (pyrexia)
- Heavy night sweets
- Rapid weight loss
- Swollen lymph nodes
- While spots on tongue, mouth, throat
- Recurrent or chronic diarrhoea
- Anxiety and depression

## Diagnosis

- Antibody /Antigen test (After18-45days after exposure of HIV
- Antibody Test (23 and 90 days after transmission)
- Nucleic acid test

## HIV screening




**HIV window period**

The time between exposure to HIV and it becomes detectable in blood is called HIV window period, during window period the test shows negative result.

## Management


- Antiretroviral therapy, its grouped into six classes
- NRTIs (nucleoside reverse trans criptase inhibitors
- NNRTIs (non-nucleoside reverse trans criptase inhibitors
- Protease inhibitors
- Fusion inhibitions
- CCR5 Antagonists
- Integrase stand transfer inhibitions

## HIV-Prevention

- Get tested for HIV
- Limit sexual partners
- Avoid sharing needless
- Stay away from drugs
- Avoid blood contact
- Practice safe sex
- Take action if you have exposed to HIV

### 2.7.2 Gonorrhoea

Gonorrhoea is a sexually transmitted disease (STD). Its caused by infection with the bacterium neisseria gonorrhoea, it tends to infect warm, moist area of the body.



## Causative organism

Nesseria Gonorrhoea

## Incubation period



XII_Nursing_Vocational_Unit 2.indd 44 1/11/2020 2:25:30 PM