2-5 daysafter exposure

Symptoms

In men

- White, yellow or green urethral discharge (pus-like)
- Pain in the testicles or scrotum
- Painful or frequent urination
- Anal discharge, itching pain, bleeding
- Itching, difficulty in swallowing swollen neck lymph nodes
- Eye pain, light sensitivity
- Eye discharge (pus-like)
- Red, swollen warm, painful joints

In women

- Painful sexual intercourse
- Fever
- Yellow or green vaginal discharge
- Vulval swelling
- Bleeding in between periods
- Heavier periods
- Bleeding after intercourse
- Vomiting, abdominal or pelvic pain Painful or frequent urination

Diagnosis

- ❖ Nucleic acid amplification test (NAAT)
- ❖ Culture
- ❖ Gram stain of urine sediment

Complications

In women

- ❖ Pelvic inflammatory disease
- ❖ Ectopic pregnancy
- ❖ Chronic pelvic pain
- ❖ Infertility

In men

- ❖ Epididymitis (in men)
- ❖ Arthritis
- ❖ Tenpsynovitis
- ❖ Dermatitis

Management

- ❖ Antibiotics (Cefltriazxone, ozithromuuin)
- ❖ Abstaining from sexual intercourse
- ❖ Eye ointment for eye infection

Gonorrhea passed to infant causes Jaint infection, blindness and also still birth, premature labour.

Preventive measures

- ❖ Safe sex practice
- ❖ Having sex with mutually monogamous, unaffected partner

XII_Nursing_Vocational_Unit 2.indd 45 1/11/2020 2:25:30 PM