2. Post operative care

- ❖ Encourage the patient to splint the incision site with hand or pillow when coughing to lessen pain and protect the site from increased intra-abdominal pressure and wound dehiscence
- ❖ Administer analgesics as ordered
- ❖ Encourage ambulation as soon as permitted
- ❖ Advise patient that difficulty in urinating is common after surgery; promote elimination to avoid discomfort, and catheterise if necessary

3. Prevention of infection

- ❖ Monitor the vital signs
- ❖ Check dressings for drainage and incision for redness and swelling
- ❖ Monitor for other signs and symptoms of infections; fever, chills, malaise, diaphoresis
- ❖ Administer prescribed antibiotics


4. Patient education on discharge

- ❖ Advise that pain and scrotal swelling may be present for 24 to 48 hours after repair of an inguinal hernia
- ❖ Apply ice intermittently
- ❖ Elevate scrotum by using scrotal support
- ❖ Take prescribed medication to relieve discomfort
- ❖ Inform that heavy lifting should be avoided for 4-6 weeks
- ❖ Athletics and extremes of extension are to be avoided for 8 to 12 weeks postoperatively

Complications

- ❖ Bowel obstruction
- ❖ Gangrene formation
- ❖ Wound dehiscence

3.2.2 Cholecystitis



Inflammation of the gall bladder

Causes

- ❖ Exact cxauses is not known
- ❖ Gall stones and kinking or twisting of bile duct

Risk factor

- ❖ Sedentary life style
- ❖ Obesity

Signs & Symptoms

- ❖ Pain on right upper quadrant/epigastric or both
- ❖ Nausea
- ❖ Vomiting
- ❖ Increased temperature
- ❖ Mild jaundice

XII_Nursing_Vocational_Unit 3.indd 58 1/11/2020 2:36:18 PM