effects evaluated as adjunctive therapy for sickle cell anaemia. Blood transfusion and folic acid therapy is administered.

5. **Aplastic anaemia:** Two methods of managements are currently employed. Bone marrow transplantation and Administration of immunosuppressive therapy with entity.

6. **Megaloblastic anaemia:** Injection B_{12} is administered, provide nutritious diet and folic acid 1 mg per day.

Nursing Management

- Offer small amount of foods at frequent intervals
- Provide iron-rich foods and vitamins.
- Teach and assist with good hygienic practices
- Good dietary habits

STUDENT'S ACTIVITY

- List the food products which you have taken whole 1 week and identify the iron nutrient in it
- Participate in WIFS (Weekly Iron and Folic acid Scheme) programme

Complication

- Mental sluggishness
- Growth retardation
- Delayed puberty related to growth retardation
- Cardiac failure related to collapse and shock resulting in death

IRON - DAILY REQUIREMENT

- Children (ages 1-10): 7 to 10 mg per day
- Women (ages 19-50): 18 mg per day
- Pregnant Women: 27 mg per day
- Lactating Women: 9 to 10 mg per day
- Men (ages 19 and older): 8 mg per day

DO YOU KNOW?

Food rich in Iron: Spinach, Beetroot, Apple, Pomegranate, Soy Beans, Nuts,Honey, Dates, Egg, Sesame Seeds, Seafood and all kind of Meats.



XII_Nursing_Vocational_Unit 3.indd 66 1/11/2020 2:36:27 PM