Nursing Management

- Pain control
- Minimize the vomiting by antiemetics
- Increased fluid intake and relaxation technique
- Provide comfortable position
- Maintain fluid intake chart (intake output)
- Advice dietary modification

3.5.2 Acute Renal Failure



Acute renal failure is a syndrome of varying causation that results in a sudden decline in renal function.

Kidneys are a pair of organs located toward lower back. One kidney is on each side of spine. They filter blood and remove toxins from body. Kidneys send toxins to your bladder. Later body removes toxins during urination. Kidney failure occurs when kidneys abruptly lose their function of filtering waste products from the blood.

Causes

- Pre–renal causes: It results from the condition that decreases renal blood flow (e.g. Hypovolemia, shock, hemorrhage, burns, diuretic therapy)
- Post–renal causes: It comes from obstruction or distruption of urine flow anywhere along the urinary tract
- Intra – renal causes: It results from injury in the renal tissue and is usually associated with intra renal ischemia, toxins immunologic process, systematic and vascular disorders

Signs and Symptoms

- Decreased tissue turger, dryness of mucous membrane, weight loss, hypotension, oliguria
- Difficulty in voiding and changes in urine flow
- Fever and skin rashes, edema
- Changes in urine volume, increases blood urea, creatinine, uric acid, potassium levels etc

Investigations

- Urine analysis: Reveals protenuria, haematuria
- Raise of Serum creatinine in blood
- Urine culture
- Renal ultrasonography
- Renal biopsy

Management

- Correct any reversible cause of acute renal failure E.g: Improve renal perfusion maximize cardiac output, surgical release of obstruction
- Be alert for fluid excess or deficit
- Monitor the signs of hypovolemia or hypervolemia
- Monitor Intake and output chart
- Weigh the patient daily

XII_Nursing_Vocational_Unit 3.indd 71 1/11/2020 2:36:30 PM