- Auscultate bowel sounds.

❖ Family education:

- Develop an interpersonal relationship with the family.
- Provide frequent update information on patient condition.
- Involve the relatives in routine care.
- Provide comfortable physical environment.
- Teach family to report any unusual symptoms.

3.7 MUSCULO SKELETAL

Osteoarthritis

Osteoporosis

3.7.1 Osteoarthritis

Osteoarthritis is a very common chronic disorder involving the joints. It is a degenerative change in the joints. The degenerative changes take place because the rubbing of the joint surfaces causes a wearing away and disintegration of the tissues.

Causes and Risk factors

Age: people above the age of 45 have the risk for developing osteoarthritis. It is, however, most commonly found in people over age of 65.


Gender: This disease of osteoarthritis is more common in women, particularly after the age of 55

Obesity: This can be one of the causes for osteoarthritis as every kilogram puts three extra kilogram of pressure on knees

Injury : Joint injuries are an increased risk of developing osteoarthritis

Weakness of muscles: Weak thigh muscles lead to osteoarthritis knee joint pain.

Stages of knee Osteoarthritis



Signs & Symptoms

❖ Knee pain
❖ Swelling
❖ Joint stiffness
❖ Loss of movement

Investigation

❖ X–ray of affected joints
❖ Serum CRP
❖ Serum ESR, RA factor

Management

Medical management:

❖ Analgesics
❖ Exercise to strengthen muscles

Surgical management:

❖ Arthodesis (practised earlier)

XII_Nursing_Vocational_Unit 3.indd 77 1/11/2020 2:36:34 PM