Medication:

- Oral hypoglycaemic agents for patient where NIDDM do not achieve glucose control with diet and exercise only
- Insulin therapy for patients with IDM who require replacement. (May also be used for NIDDM when unresponsive to diet, exercise and oral hypoglycaemic agent therapy. Hypoglycemic may result, as well as rebound hyperglycaemic effect
- Demonstrate and explain thoroughly the procedure for insulin self-injection
- Help [patient to master technique by taking a step-by-step approach
- Allow patient time-to-time handle insulin and syringe to become familiar with the equipment
- Teach self-injection first to alleviate fear from injection
- Instruct patient in filling the syringe when he or she expresses confidence in self-injection procedure
- Review dosage and time injections in relation to meals activity, and bedtime based on patient's individualized insulin regimen

Health Education:

❖ Preventing injury secondary to Hypoglycemia:

- Closely monitor blood glucose levels to detect hypoglycaemia.
- Asses patient for the signs and symptoms of hypoglycaemia.
- Sweating, cardiac palpitation and nervousness.
- Head ache, light-headedness,. Confusion, irritability, slurred speech, lack of co-ordination staggering gait from depression of central nervous system as glucose level progressively falls.
- Treat hypoglycaemia promptly with 10-15 gms of fast acting carbohydrates.
- Half-cup juice, 3 glucose tablets, 4 sugar cubes, 5-6 pieces of sugar candy may be taken orally.
- Encourage patient to carry a portable management for hypoglycaemia at all times.
- Encourage patients to wear an identification bracelet opr card that may assist in prompt management in a hypoglycaemia emergency.
- Identification bracelet may be obtained from Medic Alert Foundation.
- Identification card may be requested from the Indian Diabetes Association.
- Between meal snacks as well as extra food taken before exercise should be encouraged to prevent hypoglycaemia.

❖ Improving activity tolerance:

- Advice patient to asses blood glucose level before strenuous exercise.
- Advice patient that prolonged strenuous exercise may require

XII_Nursing_Vocational_Unit 3.indd 81 1/11/2020 2:36:40 PM