$$Z(K_n) = \frac{1}{4} \cdot \left\lfloor \frac{n}{2} \right\rfloor \cdot \left\lfloor \frac{n-1}{2} \right\rfloor \cdot \left\lfloor \frac{n-2}{2} \right\rfloor \cdot \left\lfloor \frac{n-3}{2} \right\rfloor$$

Zarankiewicz' conjecture, instead, provides an estimate for complete bipartite graphs with two partitions having `n` and `m` vertices respectively:

$$Z\left(K_{n,m}\right) = \left\lfloor \frac{n}{2} \right\rfloor \cdot \left\lfloor \frac{n-1}{2} \right\rfloor \cdot \left\lfloor \frac{m}{2} \right\rfloor \cdot \left\lfloor \frac{m-1}{2} \right\rfloor$$

As of today, both formulas have been proven to hold as upper bounds, meaning that the crossing number for these graphs is not larger than the value computed using the formulas, and they were not yet been disproved as lower bounds.

If we try to apply these results to the two graphs we have already introduce, we get:

$$Z(K_5) = \frac{1}{4} \cdot \left\lfloor \frac{5}{2} \right\rfloor \cdot \left\lfloor \frac{4}{2} \right\rfloor \cdot \left\lfloor \frac{3}{2} \right\rfloor \cdot \left\lfloor \frac{2}{2} \right\rfloor = 1$$

$$Z(K_{3,3}) = \left\lfloor \frac{3}{2} \right\rfloor \cdot \left\lfloor \frac{2}{2} \right\rfloor \cdot \left\lfloor \frac{3}{2} \right\rfloor \cdot \left\lfloor \frac{2}{2} \right\rfloor = 1$$

So, for K_5 and $K_{3,3}$ the expectation is consistent with what we have mentioned, and with our experience. As a matter of