```
var yScale = d3.scale.linear() .domain([0,24500]).range([0,100]);

d3.select("svg")
  .selectAll("rect")
  .data([14, 68, 24500, 430, 19, 1000, 5555])
  .enter()
  .append("rect")
  .attr("width", 10)
  .attr("height", function(d) {return yScale(d);})
  .style("fill", "blue")
  .style("stroke", "red")
  .style("stroke-width", "1px")
  .style("opacity", .25)
  .attr("x", function(d,i) {return i * 10;})
  .attr("y", function(d) {return 100 - yScale(d);});
```

When you deal with such widely diverging values, it often makes more sense to use a polylinear scale. A polylinear scale is a linear scale with multiple points in the domain and range. Let's suppose that for our dataset, we're particularly interested in values between 1 and 100, while recognizing that sometimes we get interesting values between 100 and 1000, and occasionally we get outliers that can be quite large. We could express this in a polylinear scale as follows:

```
var yScale =
d3.scale.linear().domain([0,100,1000,24500]).range([0,50,75,100]);
```

The previous draw code produces a different chart with this scale, as shown in figure 2.18.




Figure 2.17 A bar chart drawn using a linear scale




Figure 2.18 The same bar chart from figure 2.17 drawn with a polylinear scale