```
    m /= m.sum()                                    6
    return m
```

- ***1*** Calculates the support spacing value
- ***2*** Calculates the index value of the observed reward in the support
- ***3*** Rounds and clips the value to make sure it is a valid index value for the support
- ***4*** Starting from the immediate left neighbor, steals part of its probability
- ***5*** Starting from the immediate right neighbor, steals part of its probability
- ***6*** Divides by the sum to make sure it sums to 1

Let's walk through the mechanics of this to see how it works. We start with a uniform prior distribution:

```
>>> probs
array([0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784, 0.01960784, 0.01960784, 0.01960784, 0.01960784,
       0.01960784])
```

You can see that each support has a probability of about 0.02. We observe $r_t = -1$, and we calculate $b_j \approx 22$. We then find the nearest left and right neighbors, denoted m_l and m_r, to be indices 21 and 23, respectively. We multiply m_l by γ^j, where j is a value that we increment by 1 starting at 1, so we get a sequence of exponentially decreasing gammas: $\gamma^1, \gamma^2, \ldots \gamma^j$. Remember, gamma must be a value between 0 and 1, so the sequence of gammas will be 0.5, 0.25, 0.125, 0.0625 if $\gamma = 0.5$. So at first we take 0.5 * 0.02 = 0.01 from the left and right neighbors and add it to the existing probability at $b_j = 22$, which is also 0.02. So the probability at $b_j = 22$ will become 0.01 + 0.01 + 0.02 = 0.04.