67. A biodiversity hope spot is an area of ocean that is under special protection because of its wildlife and significant under water habitat. They are places where human activities and impacts will be minimised, the unique and vital ecosystems within the Hope Spots will have a chance to recover and to flourish. Biodiversity hope spot program is overseen by Mission Blue in partnership with IUCN.

68. A food web illustrates the numbers of each organism which are eaten by others, but food chain does not. Food chains can be found within the Communities of an ecosystem as communities consist of different species to feed upon.

69. Low temperature and anaerobiosis, in taiga, inhibit decomposition resulting in a build-up of organic materials soils become acidic. In tropical rain forests, the nutrients are quickly released due to abundance of heat and air. But these nutrients are either quickly absorbed by dense shrubs on ground or washed away by rain because of which soils are deprived of nutrients ^ soils become acidic.

70. All food pyramids, including pyramids of energy, biomass and number, do not consider the same species belonging to two or more trophic levels.

Food pyramids assume a simple food chain, something that seldom exists in nature; it does not accommodate a food web which considers the same species belonging to two or more trophic levels.

71.

- In freshwater the average concentration of dissolved oxygen is ten parts per million by weight.
- This is 150 times lower than the concentration of oxygen in an equivalent volume of air. Oxygen enters the aquatic ecosystem through the air-wa- ter interface and by the photosynthetic activities of aquatic plants.
- Dissolved oxygen escapes the water body through the air-water interface and respiration of organisms (fish, decomposers, zooplankton, etc.).
- The amount of dissolved oxygen retained in water is also influenced by temperature.