- Oxygen is less soluble in warm water → more dissolved oxygen would escape into atmosphere. Warm water also enhances decomposer activity → consumes dissolved oxygen reducing amount of oxygen available for other species.
- When the dissolved oxygen level falls below 3-5 ppm, many aquatic organisms are likely to die leading to reduced diversity.

72. Wetlands are areas of marsh or peatland with water that is static or flowing, fresh, brackish or saline, including areas of marine water the depth of which at low tide does not exceed 6 m.

Wetlands are transition zones (ecotone) between terrestrial and aquatic ecosystems.

According to the Directory of Asian Wetlands (1989), wetlands occupy 18.4% of the country's area (excluding rivers), of which 70% are under paddy cultivation.

Recent estimates of the total area of wetlands in India include the following:

Area under paddy cultivation	40,990,000 ha
Area suitable for fish culture	
(a) freshwater 1,600,000 ha	
(b) brackish water 2,000,000 ha	3,600,000 ha
Area of capture fisheries	2,900,000 ha
Mangroves (after Blasco 1977)	356,000 ha
Estuaries	3,900,000 ha
Back waters	3,540,000 ha
Man-made impoundments	3,000,000 ha
Rivers including main tributaries	28,999 km
Canals and irrigation channels	113,000 km
Total area of wetlands (Excluding rivers)	58,286,000 ha

73.

- The extent of sulphur is the major difference between Bharat Stage IV and Bharat Stage VI norms.
- BS-IV fuels contain 50 parts per million (ppm) sulphur; the BS-VI grade fuel only has 10 ppm sulphur.