- Black carbon particles (a component of soot) originating from combustion processes have been known for some time to absorb sunlight and warm the atmosphere, and pollution controls have been put into place to reduce their emissions and their effects.
- Brown carbon has attracted interest as a possible cause of climate change. This class of organic carbon, known for its brownish colour, is absorbed strongly in the ultraviolet wavelengths and less significantly going into the visible wavelengths. Types of brown carbon include tar materials from smouldering fires or coal combustion, breakdown products from biomass burning, a mixture of organic compounds emitted from soil, and volatile organic compounds given off by vegetation.

91. Effect of Nano Particles on stratospheric temperature
    - NPs in the troposphere interact with molecular hydrogen accidentally released from hydrogen fuel cells and other sources.
    - Molecular hydrogen along with NPs moves up to the stratosphere, resulting in the abundance of water vapour in the stratosphere.
    - This will cause stratospheric cooling due to the formation of Stratospheric clouds (mostly ice crystals).
    - Stratospheric clouds destroy ozone.
92. UN-REDD:
    - The United Nations Programme on Reducing Emissions from Deforestation and Forest Degradation.
    - Headquarters: Geneva, Switzerland.
    - Membership: 64 Partner Countries.
    - It is a multilateral collaborative programme of the Food and Agriculture Organization of the United Nations (FAO), United Nations Development Programme (UNDP) and United Nations Environment Programme (UNEP).
    - It was created in 2008 in response to the UNFCCC decisions on the Bali Action Plan and REDD at COP-13. It partners with developing countries to support them in establishing the technical capacities needed to implement REDD+ and meet UNFCCC requirements for REDD+ results- based payments.