examples of the Hot desert and mid-latitude deserts can be located in Asia.

Statement 4: CORRECT—The warm Temperate western margin climate is also called as Mediterranean climate. Wet winters and dry summers are typical of the Mediterranean climatic region in West Asia, where precipitation is associated with the winter activity of the polar front. That polar-front activity, accompanied by maximum precipitation, occurs in the spring in the interior parts of the West Asian highlands. Summer and winter precipitation merges in some parts of Asia.

53. The distribution pattern of precipitation throughout the year is varied. Relatively uniform moisture is characteristic of the Asian equatorial zone. Maximum summer precipitation and minimum winter precipitation are the rule in the subequatorial zones and in other regions with monsoon climates, as well as in those areas where there is summer movement of the fronts—the polar front in the mountains of southern Siberia and the Arctic front in the subarctic regions. Wet winters and dry summers are typical of the Mediterranean climatic region in West Asia, where precipitation is associated with the winter activity of the polar front. That polar-front activity, accompanied by maximum precipitation, occurs in the spring in the interior parts of the West Asian highlands. Summer and winter precipitation merges in some parts of Asia. In the Kolkhida area of Georgia east of the Black Sea, the summer rains —brought by the northwesterly Atlantic air currents—merge with the cyclonic Mediterranean winter rains. In some areas of Japan, Korea, and eastern China, there is uniform precipitation when, in addition to the summer monsoon, the winter monsoon brings moisture.
54. The tundra climate, associated with the cold, treeless plains of the Arctic lowlands of Asia. The cold, sharply continental climate of eastern Siberia, the cold, moderately humid western Siberian climate, the humid, subtropical climate associated with the Kolkhida region, the desert climate of the temperate zone.
55. Statement 1: CORRECT—In the Severnaya Zemlya archipelago, off the Arctic coast, thawing begins in May and frosts begin in August, although in some years frosts may occur at night