throughout the short summer.
Statement 2: CORRECT—Tundra extends over large areas of the Chersky, Verkhoyansk, and Kamchatka mountain ranges.
Statement 3: INCORRECT—The taiga has a thin undergrowth of cranberries and bilberries, and there are numerous extensive peat bogs.

A peat bog is a type of wetland whose soft, spongy ground is composed largely of living and decaying Sphagnum moss. Decayed, compacted moss is known as peat, which can be harvested to use for fuel or as a soil additive. Peat bogs are found throughout the world where cool temperatures and adequate rainfall prevail.

56. Lake Baikal is a rift lake located in southern Siberia, Russia.
Lake Balkhash is one of the largest lakes in Asia and 15th largest in the world. It is located in Central Asia in southeastern Kazakhstan
Tso Moriri or Lake Moriri or “Mountain Lake”, is a lake in the Changthang Plateau (literally: northern plains) in Ladakh in Northern India.
Poyang Lake located in Jiangxi Province, is the largest freshwater lake in China.

57. About two-thirds of Japan’s total area is forested, whereas much of China is deforested (roughly one- seventh of China is under forest cover); sizable tracts remain untouched, however, in the remote rugged regions of China, and many small areas have been reforested. One reason for that difference between the two countries is the traditional respect the Japanese have had for their scenery and, consequently, the strict forestry regulations that have been promulgated and enforced. In addition, the rugged topography and inaccessibility of many of Japan’s forested areas have limited the possibilities for economic exploitation. The best examples of East Asian forest, therefore, are found in Japan, such as in the Kii Peninsula of Honshu.

58. The wettest parts of peninsular India (such as the Western Ghats) and of Southeast Asia have magnificent tropical forests noteworthy for the variety of their plant life. A significant feature of South Asian vegetation is the family Dipterocarpaceae (yielding aromatic oils