Paraguay, parts of Bolivia, Brazil, Argentina, and Chile) and in the mid-level elevations of the Andes.

30. Areas where average annual temperatures are less than 50 °F (10 °C) are characterised as cold climates. Those occur in the southernmost parts of Argentina and Chile and in the high Andes above about 11,500 feet (3500 metres). Mean temperatures are relatively low throughout the year, but daily variations are wide. There is a marked difference in humidity between the northern and southern parts of the upper Andean zone. In Colombia and Ecuador the climate at such elevations is cool and damp. Temperatures, always low, may on the average vary daily from 54 °F (12 °C) during the daytime to 28 °F (-2 °C) at night. Precipitation generally is high and well distributed throughout much of the year, although most of the Ecuadorian Andes have a dry period from June to August. Clouds and mist are dense in much of the region, and sunlight penetrates only for short periods. From central Peru to Bolivia and Chile, temperatures are still lower. Near Lake Titicaca, the average annual temperature is only about 34 to 36 °F (1 to 2 °C).
31. Tropical savannas are found mostly in the Llanos of Venezuela and north-eastern Colombia. Those vast plains are covered with grasses and sedges, but tree clusters (mainly palms) also grow, especially along streams.

    The flat or softly rolling plains called Pampas, which constitute the greater part of eastern Argentina, are covered with grasses.

    The ***campo rupestre*** (“rupestrian grassland”) is a discontinuous montane subtropical ecoregion occurring across three different biomes in Brazil: Cerrado, Atlantic Forest and Caatinga.
32. South Brazilian forests:

    Those parklike forests, sometimes very dense but interspersed with savanna, occupy vast expanses from the border of the Amazonian rainforest to the marshes of the upper Paraguay River. The typical landscape is a grassland strewn with smaller trees. In effect, it includes a mosaic of associations, from hygrophilous (living or growing in moist places) to xerophilous (adapted to dry conditions) forests and even desert.