soyabean. Some of the most important rice-growing regions are Assam, West Bengal, coastal regions of Odisha, Andhra Pradesh, Telangana, Tamil Nadu, Kerala and Maharashtra, particularly the (Konkan coast) along with Uttar Pradesh and Bihar.

Statement III: CORRECT—In between the rabi and the kharif seasons, there is a short season during the summer months known as the Zaid season. Some of the crops produced during 'zaid' are watermelon, muskmelon, cucumber, of canal irrigation and tube- wells have made it possible to grow rice in areas of less rainfall such as Punjab, Haryana and western Uttar Pradesh and parts of Rajasthan.

57. Statement I: CORRECT—Sugarcane: It is a tropical as well as a subtropical crop. It grows well in hot and humid climate with a temperature of 21°C to 27°C and an annual rainfall between 75 to 100 cm. Irrigation is required in the regions of low rainfall. It can be grown on a variety of soils and needs manual labour from sowing to harvesting.

India is the second largest producer of sugarcane only after Brazil. It is the main source of sugar, gur (jaggary), khandsari and molasses. The major sugar- cane-producing states are Uttar Pradesh, Maharashtra, Karnataka, Tamil Nadu, Andhra Pradesh, Telangana, Bihar, Punjab and Haryana.

Statement II: CORRECT—Oil Seeds: In 2014 India was the second largest producer of groundnut in the world after China. In rapeseed production India was third largest producer in the world after Canada and China in 2014.

Different oil seeds are grown covering approximately 12% of the total cropped area of the country. Main oil-seeds produced in India are groundnut, mustard, and coconut, sesamum (til), soyabean, castor seeds, cotton seeds, linseed and sunflower. Most of these are edible and used as cooking mediums. However, some of these are also used as raw material in the production of soap, cosmetics and ointments.

58. Statement I: CORRECT—Irrigation is a major determinant of agricultural productivity. Indian agriculture has been constrained by limited irrigation and only about 40.6% of arable land is irrigated. The remaining 60% continues to depend on rainfall. The