The most important earthquake belt is the Circum- Pacific Belt. The mid-world mountain belt (Alpine Belt) extends parallel to the equator from Mexico across the Atlantic Ocean, the Mediterranean Sea from Alpine-Caucasus ranges to the Caspian, Himalayan mountains and the adjoining lands also experiences many major earthquakes. Hence is not an exception. **Wadati Benioff** zone is a zone of subduction along which earthquakes are common. A Wadati-Benioff zone is a zone of seismicity corresponding with the down-going slab in a subduction zone (Convergent Boundary). Differential motion along the zone produces numerous earthquakes. Shallow focus earthquakes (most common at submarine ridges. Hardly felt). Intermediate focus earthquakes (somewhat severe). Deep focus earthquakes (Occurs at trenches - convergent boundary and are very powerful. Japan lies along trench line. Hence it faces devastating earthquakes).

35. Granite is an intrusive volcanic rock formed by slow cooling of crystals of lava which allows formation larger grains. Basalt is a plutonic rock formed by rapid cooling of crystal of lava results in fine grained rocks.
36. A gully is a landform created by running water. Gullies resemble large ditches or small valleys but are metres to tens of metres in depth and width. A ravine is a landform narrower than a canyon and is often the product of stream cutting erosion. Ravines are typically classified as larger in scale than gullies, although smaller than valleys.
37. The wind or Aeolian erosion takes place in the following ways, viz. deflation, abrasion, and attrition. Deflation is removing, lifting and carrying away dry, unsorted dust particles by winds. It causes depressions known as blow outs or deflation blowouts.
38. The wind or Aeolian erosion takes place in the following ways, viz. deflation, abrasion, and attrition. Deflation is removing, lifting and carrying away dry, unsorted dust particles by winds. It causes depressions known as blow outs or deflation blowouts.
39. Laccolith formed by intrusive volcanism and Mushroom Rocks formed due to Aeolian Erosion are most common mushroom shaped rocks formed by geomorphic processes. Though mushroom rocks can be formed by glacial erosion, such rocks