experimenters to start constructing bar detectors. By 1975 it will be clear he was not seeing waves. [Ch. 10]

Braginsky discovers evidence that there will be a quantum limit on the sensitivities of gravitational-wave detectors. [Ch. 10]

1970 Bardeen shows that the accretion of gas is likely to make typical black holes in our Universe spin very rapidly. [Ch. 9]

Price, building on work of Penrose, Novikov, and Chase, de la Cruz, and Israel, shows that black holes lose their hair by radiating it away, and he proves that anything which can be radiated will be radiated away completely. [Ch. 7]

Hawking formulates the concept of a black hole's absolute horizon and proves that the surface areas of absolute horizons always increase. [Ch. 12]

Giacconi's team constructs Uhuru, the first X-ray detector on a satellite; it is launched into orbit. [Ch. 8]

1971 Combined X-ray, radio-wave, and optical observations begin to bring strong evidence that Cygnus X-1 is a black hole orbiting a normal star. [Ch. 8]

Weiss at MIT and Forward at Hughes pioneer interferometric detectors for gravitational waves. [Ch. 10]

Rees proposes that a radio galaxy's giant lobes are powered by jets that shoot out of the galaxy's core. [Ch. 9]

Hanni and Ruffini formulate the concept of surface charge on a horizon, a foundation for the membrane paradigm. [Ch. 11]

Press discovers that black holes can pulsate. [Ch. 7]

Zel'dovich speculates that spinning black holes radiate, and Zel'dovich and Starobinsky use the laws of quantum fields in curved spacetime to justify Zel'dovich's speculation. [Ch. 12]

Hawking points out that tiny "primordial" black holes might have been created in the big bang. [Ch. 12]