




Ray-Ban Glasses with FB tech

Ray-Ban Stories powered by Facebook's technology that can record videos upto 30secs <https://digit.in/oct21-11>




EU rule forces USB-C chargers

Manufacturers will be forced to create a universal charging solution for 6 small electronic devices <https://digit.in/oct21-12>

1000 WORDS


Bessel Function


In 1817, when German astronomer Friedrich Bessel was working on a solution of one of the Kepler's equations on planetary motion After Bessel published his findings, other scientists found that the functions appeared in mathematical descriptions of many physical phenomena, including the flow of heat or electricity in a solid cylinder.




Thomae's function on [0,1]

The only function that has a number of interesting names is Thomae's function, named after Carl Thomae, a student of Reimann. The names given to this function are: The popcorn function, the raindrop function, the countable cloud function, the modified Dirichlet function, the ruler function, the Riemann function and the stars over Babylon function.

Advanced Calculus




The base of every technical field is calculus, including engineering, computer science, and architecture. Even biologists use advanced calculus to determine the growth rate of bacteria culture when the surroundings like temperature and food are changed. Life would be boring without calculus.



Mandelbrot Set

Mandelbrot set is a fractal and is similar to the Julia set. The Mandelbrot set has become famous outside mathematics too, for its appealing beauty and its complex structure arising from simple mathematical concepts.



Gamma function

The gamma function is a factorisation of complex numbers, which are a series of numbers represented by the series $a+bi$, where a and b are real numbers and i is the imaginary unit. Except for non positive integers, the gamma function is defined for all complex numbers.

Euler's polyhedron formula

The number of vertices V , faces F , and edges E in a convex 3-dimensional polyhedron, satisfy $V + F - E = 2$, Euler's formula may look simple but when applied can create beautiful shapes with tons of faces, vertices, and edges, imagine the potential.

